

FOOD  
INNOVATION  
ASIA  
CONFERENCE  
2016

# PROCEEDINGS FOOD RESEARCH AND INNOVATION FOR SUSTAINABLE GLOBAL PROSPERITY

16 -18 JUNE 2016  
BITEC BANGNA, BANGKOK, THAILAND









**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

**About the Conference**

The conference will provide opportunity to meet and share experiences as well as strengthen networking among international food scientists and scientists in related fields from academia, government and food industries. The objective is to highlight significant developments in research and innovations in food science and technology with an emphasis on Food Research and Innovation for Sustainable Global Prosperity. The conference will feature a series of presentations and discussions in plenary, concurrent and poster sessions, informal gatherings, competitions and exhibitions.

- Division (A) Food Chemistry, Nutrition, and Analysis
- Division (B) Food Processing and Engineering
- Division (C) Food Product Development, Sensory, and Consumer Research
- Division (D) Food Microbiology, Food Biotechnology, Fermentation
- Division (E) Related Topics (Food Packaging, Food Safety & Quality, Food Laws & Regulations, Food Policy, etc.)
- Division (F) Industrial session

**Concurrent Events**

- o ProPak ASIA 2016
- o FoSTAT - Nestlé Quiz Bowl 2016
- o Food Innovation Contest 2016 (final round)
- o STI for Agri-Food Industry Development Framework
- o ACS Thailand section
- o FIFSTA Annual meeting
- o AIAC Annual meeting
- o Thailand Section of AOAC International meeting

**Organizers**

- o Food Science and Technology Association of Thailand (FoSTAT)
- o Agro-Industry Academic Council Association (AIAC)
- o Faculty of Agro-Industry , Chiang Mai University
- o School of Agriculture Technology, Walailak University
- o National Science Technology and Innovation Policy Office (STI)
- o Thailand Section of AOAC International

**Supported by**

- o Bangkok Exhibition Services Ltd. (BES)
- o The Federation of the Institute of Food Science and Technology in ASEAN (FIFSTA)



**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
16–18 June 2016, BITEC Bangna, Bangkok, Thailand

**List of Food Innovation Asia Conference 2016 Committee**

**Organizing Committee**

- o Professor Dr.Pavinee Chinachoti
- o Associate Professor Dr.Tanaboon Sajjaanantakul
- o Associate Professor Dr.Anadi Nitithamyong
- o Associate Professor Dr.Saiwarun Chaiwanichsiri
- o Assistant Professor Dr.Charin Techapun
- o Assistant Professor Dr.Suwit Wuthisuthimethavee

**Scientific Committee**

**Food Chemistry, Nutrition, and Analysis**

- o Associate Professor Dr.Manat Chaijan
- o Associate Professor Dr.Patcharin Raviyan
- o Assistant Professor Dr.Worawan Panpipat
- o Dr.Akkasit Jongjareonrak
- o Assistant Professor Dr.Panida Rattanapitigorn
- o Dr.Prodpran Thakeow Angeli
- o Dr.Sutee Wangtueai
- o Dr.Suthasinee Yarnpakdee
- o Lecturer Khamtorn Pudtikajorn
- o Lecturer Wachira Jirattananarangsri
- o Assistant Professor Dr.Supranee Manurakchinakorn
- o Assistant Professor Dr.Thunnop Laokuldilok

**Food Processing and Engineering**

- o Associate Professor Dr.Aphirak Phianmongkhol
- o Associate Professor Dr.Noppol Leksawasdi
- o Assistant Professor Dr.Pattavara Pathomrungsyounggul
- o Assistant Professor Dr.Rattana Muangrat
- o Assistant Professor Dr.Somchai Jomduang
- o Assistant Professor Dr.Srisuwan Naruenartwongsakul
- o Assistant Professor Dr.Yongyut Chalermchat
- o Assistant Professor Kongdej Limpaboon
- o Dr.Akkasit Jongjareonrak
- o Dr.Pilairuk Intipunya
- o Dr.Supavej Maniyom
- o Dr.Tanong Aewsiri
- o Lecturer Ravis Tasakorn
- o Dr.M.L.Yasinee Chakrabandhu



**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

o Dr.Visaka Anantawat

**Food Product Development, Sensory, and Consumer Research**

- o Professor Dr.Pairote Wiriyaacharee
- o Assistant Professor Dr.Narumol Matan
- o Associate Professor Dr.Niramom Utama-ang
- o Assistant Professor Dr.Sujinda Sriwattana
- o Assistant Professor Dr.Suthat Surawang
- o Assistant Professor Dr.Yuthana Phimolsiripol
- o Lecturer Israpong Pongsirikul
- o Lecturer Kanyarat Suthapakti
- o Lecturer Suwanna Decharatanangkoon
- o Lecturer Vinyoo Sakdatorn
- o Lecturer Wiwat Wattanatchariya
- o Dr.Piyawan Simapaisarn
- o Dr.Tanong Aewsiri

**Food Microbiology, Food Biotechnology and Fermentation**

- o Associate Professor Dr.Wanna Choorit
- o Assistant Professor Dr.Ampin Kuntiya
- o Assistant Professor Dr.Charin Techapun
- o Assistant Professor Dr.Chartchai Khanongnuch
- o Assistant Professor Dr.Nisa Saelee
- o Assistant Professor Dr.Phisit Seesuriyachan
- o Assistant Professor Dr.Prasert Hanmoungjai
- o Assistant Professor Dr.Sasitorn Wongruong
- o Dr.Churairat Moukamnerd
- o Dr.Pijug Summpunn
- o Dr.Pornpimol Ouichanpagdee
- o Dr.Ronachai Pratanaphon
- o Dr.Siriwat Jinsiriwanit
- o Dr.Sukhuntha Osiriphun
- o Dr.Suphawatt Sinsuwongwat
- o Assistant Professor Dr.Thanongsak Chaiyaso
- o Assistant Professor Dr.Tri Indrarini Wirjantoro
- o Assistant Professor Dr.Narumol Matan
- o Dr.Sasitorn Baipong



**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

**Related Food Topics (Food Packaging, Food Service, Marketing & Management, etc.)**

- o Associate Professor Dr.Manat Chaijan
- o Associate Professor Dr.Pornchai Rachtanapun
- o Assistant Professor Dr.Pichaya Poonlarp
- o Assistant Professor Dr.Jurmkwon Sangsuwan
- o Dr.Kittisak Jantanasakulwong
- o Dr.Linda Thiraphattaraphun
- o Dr.Sutthira Sutthasupa
- o Lecturer Somchai Wongsuriyasak
- o Assistant Professor Dr.Worawan Panpipat
- o Dr.Sorraya Khiewnavawongsa

**Poster Presentation Competition**

- |   |                           |
|---|---------------------------|
| o Associate Professor Dr.Jirawat Yongsawatdigul | Chair                     |
| o Associate Professor Dr.Sappasith Klomklao     | Committee                 |
| o Associate Professor Dr.Weerachet Jittanit     | Committee                 |
| o Dr.Inthawoot Suppavorasatit                   | Committee                 |
| o Dr.Sukhuntha Osiriphun                        | Committee                 |
| o Dr.Sitthipong Nalinanon                       | Committee                 |
| o Associate Professor Dr.Manat Chaijan          | Committee and Secretariat |
| o Assistant Professor Dr.Worawan Panpipat       | Secretariat               |

**Oral Presentation Competition Committee**

- |  |                           |
|--|---------------------------|
| o Associate Professor Dr.Sanguansri Charoenrein    | Chair                     |
| o Associate Professor Dr.Wanna Choorit             | Committee                 |
| o Assistant Professor Dr.Soisuda Pornpukdeewattana | Committee                 |
| o Dr.Sasitorn Baipong                              | Committee                 |
| o Assistant Professor Dr.Ratchanee Charoen         | Committee and Secretariat |







**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

<b>Division D: Food Microbiology, Food Biotechnology and Fermentation</b>			
<b>No./Ref.</b>		<b>Title</b>	<b>Page</b>
1	OD47	Xylanase production by <i>Aspergillus niger</i> in solid state fermentation	85
2	OD51	Enzymatic production of xylooligosaccharides from alkali-soluble hemicellulose of defatted rice bran	92

<b>Division E: Related Topics (Food Packaging, Food Safety &amp; Quality, Food Laws &amp; Regulations, Food Policy, etc.)</b>			
<b>No./Ref.</b>		<b>Title</b>	<b>Page</b>
1	OE29	Challenges and difficulties in implementing food safety management system towards BRC certified food productions in Thailand	100
2	OE33	Immune enhancing effects of fermented Noni products <i>in vivo</i> and <i>in vitro</i>	107
3	OE206	Comparisons between conventional, ultrasonic-assisted, microwave-assisted and combination methods for extraction of bioactive compounds from stink bean ( <i>Parkia speciosa</i> Hassk.) pod	116

**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
16–18 June 2016, BITEC Bangna, Bangkok, Thailand

<b>Session : Poster Presentation</b>			
<b>Division A: Food Chemistry, Nutrition and Analysis</b>			
<b>No./Ref.</b>		<b>Title</b>	<b>Page</b>
1	P12	Effect of amino acid and storage temperature on kinetics of garlic discoloration	124
2	P13	Chemical qualities (pH values, TSS, total polyphenol content) in Pennywort juices producing by home cooking methods	131
3	P17	Use of the combined phase partitioning systems for lipase recovery from Pacific white shrimp ( <i>Litopenaeus vannamei</i> ) hepatopancreas	136
4	P30	In vitro studies on the antioxidant property and inhibition of angiotensin converting enzyme (ACE) from mango ( <i>Mangifera indica</i> L.) extract	143
5	P43	Nutritional quality of modified soy milk and feasibility of coconut milk replacement by modified soy milk in a dessert (ka-nhom-krok) in home cooking	150
6	P50	Influence of quinoa variety, soaking and salt concentration on percentage of germination and antioxidant activity of quinoa	157
7	P54	Production and quality of sterilized liquid non-dairy creamer from refined rice bran oil	163
8	P60	Antidiabetic activity of anthocyanins extract of black purple rice ( <i>Oryza sativa</i> L.)	172
9	P66	Antioxidant activity of Glochidion sphaerogymum extract in o/w emulsion system at various pHs and storage temperatures	181
10	P71	Development of Gymnema indorum based food controlling blood sugar	190
11	P76	Effects of three hydrocolloids on physicochemical and starch digestive properties of frozen cooked waxy rice	197
12	P77	Efficacy of plants extract, Macleaya cordata, as feed attractant in diet for Nile tilapia ( <i>Oreochromis niloticus</i> )	208
13	P79	Red aril and yellow pulp from Gac fruit ( <i>Momordica cochinchinensis</i> Spreng) as a source of biologically active ingredients for functional drink: Effect of formulation	215
14	P80	Changes of chemical properties and functional compounds during the germination of various brown rice in Mekong delta, Viet Nam	223
15	P88	Effect of rice bran flour on physicochemical properties and sensory evaluation of reduced fat pate	231
16	P94	Instant organic rice bran milk: Production and its primarily quality attributes	239

**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

17	P95	Enhance cooked shrimp color by astaxanthin and capsanthin in white leg shrimp ( <i>Litopenaeus vannamei</i> ) diets	248
18	P103	Cooking quality and anthocyanin stability as affected by cooking methods and storage time of sangyod brown rice	255
19	P106	Effects of astaxanthin and capsanthin on red tilapia ( <i>Oreochromis niloticus</i> ) skin colour intensity and growth performance	264
20	P112	Development of texture-modified food recipes for Thai elderly with swallowing difficulty	270
21	P127	Increasing manganese deposition in Asian seabass ( <i>Lates calcarifer</i> ) from dietary manganese amino acid complex	277
22	P128	Influence of preparation methods on color, antioxidant activity and physicochemical properties of purple sweet potato flour	283
23	P135	Effects of alkali solutions on the production of alkalized quail eggs	291
24	P136	Production of alkalized quail eggs from alkali solution with natural extracts	297
25	P139	Antioxidant activity and phenolic, flavonoid content of some Thai indigenous plant extracts	302
26	P141	Effect of different processing operations on total phenolic compounds of landrace rice varieties	309
27	P153	Improving hybrid catfish ( <i>Clarias macrocephalus</i> x <i>Clarias gariepinus</i> ) fillet color by lutein and organic acid in diet	317
28	P160	Enhance zinc deposition in Asian Seabass ( <i>Lates calcarifer</i> ) from dietary zinc amino acid complex	323
29	P162	Enhance calcium and phosphorus deposition in hybrid catfish ( <i>Clarias macrocephalus</i> x <i>Clarias gariepinus</i> ) by phytase enzyme in diet	329
30	P175	Chemical properties, <i>in vitro</i> starch digestibility and enzymatic inhibitory activity of cooked colored rice	336
31	P190	Guar meal digestibility in tilapia ( <i>Oreochromis niloticus</i> ) and effect on growth performance	344
32	P194	Phenolic compounds and antioxidant capacity of nine genotypes of Thai rice ( <i>Oryza sativa</i> L.) from neung rai neung san planting technique	349
33	P198	Application of near-infrared spectroscopy for determination of xanthones in mangosteen rind	358
34	P225	Essential fatty acid contents of pacific white shrimp ( <i>Litopenaeus vannamei</i> ) from krill meal in the diets of shrimp	363
35	P234	Effect of pre-treatments on physical and chemical attributes of corn nut	369
36	P235	Comparison of fresh and frozen squeezed pulp of a	377

**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

		ripe palmyra palm on colour, phenolics and carotenoid and antioxidant activity	
37	P243	Alkaline extraction and characterization of rice bran protein concentrate from commercial defatted rice bran flour	384
38	P244	Product development of reduced sugar gummy jelly from <i>Tremella fuciformis</i> (white jelly fungus)	391
39	P246	Biogenic amine determination in fresh-water fish fish-sauce by HPLC with pre-column derivatization	398
40	P11	Metabolomics signature of liquid milk and yoghurt revealed by <sup>1</sup> H-NMR technique	404
41	P36	The anti-oxidant and anti-microbial properties of young leaves of Ma-kiang ( <i>Cleistocalyx nervosum</i> var. <i>paniala</i> ) as a novel source of bioactive compounds	411
42	P65	Improving boiled shrimp color by astaxanthin in diet for white shrimp ( <i>Litopenaeus vannamei</i> )	418
43	P101	Sacha inchi protein hydrolysates: physical and functional properties	424
44	P161	Antioxidant activities of seeds and leaves of <i>Leucaena leucocephala</i>	435



**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

<b>Session : Poster Presentation</b>			
<b>Division C: Food Product Development, Sensory and Consumer Research</b>			
<b>No./Ref.</b>		<b>Title</b>	<b>Page</b>
1	P28	Formulation of khanom-piak-poon with <i>Stevia rebaudiana</i> bertonii as natural sweeteners	529
2	P52	Qualities of gluten free sponge cake substituted wheat flour with tapioca starch	534
3	P56	Sensory lexicon development for oyster sauce	540
4	P82	Development of tofu from various beans	548
5	P83	Development of functional milk drink from colored rice bran	556
6	P85	Development of salad dressing mixed with Jerusalem artichoke flour	564
7	P113	Adding values to fruit juice and mulberry's by-product through alcoholic fermentation	572
8	P117	Development of whole wheat bread from wheat-jackfruit seed mixed flour	580
9	P132	Effect of rice flour type on the quality of flour-based part of frozen kanom krok	587
10	P142	Noodles fortification with gac powder: sensory evaluation and effect of boiling and stir-frying on beta-carotene content	594
11	P189	Shelf life of pasteurized soursop ( <i>Annona muricata</i> L.) leaf tea products	601
12	P199	The development of high dietary fiber crispy roti	610
13	P223	Substitution of job's tears flour on the quality of gluten-free bread	616
14	P236	Rheological characterization of pre-emulsified oil and fish sausage batter as affect by soy protein isolate and sodium caseinate	624
15	P163	The formulation of dried noodle made of mocaf, wheat flour, protein concentrates of cowpea ( <i>Vigna unguiculata</i> , L) and porang ( <i>Amorphophallus onchophyllus</i> ) flour	632

**The 18<sup>th</sup> Food Innovation Asia Conference 2016 (FIAC 2016)**  
**16–18 June 2016, BITEC Bangna, Bangkok, Thailand**

<b>Session : Poster Presentation</b>			
<b>Division D: Food Microbiology, Food Biotechnology and Fermentation</b>			
<b>No./Ref.</b>		<b>Title</b>	<b>Page</b>
1	P19	Effect of sodium lactate on inhibition of <i>Clostridium perfringens</i> spores in sous-vide model (SVM) broth	642
2	P20	Monitoring survival of <i>Salmonella</i> and <i>Listeria</i> in semi-dry nham during storage	648
3	P32	Green synthesis and characterization of silver nanoparticles using Nang Lae pineapple ( <i>Ananas Comosus L. Merr</i> ) peel extract	655
4	P40	Antimicrobial activity of isothiocyanate from Thai cruciferous vegetables	661
5	P59	<i>In vitro</i> susceptibility of <i>Penicillium</i> spp. to vapor phase-ethanol and vinegar	669
6	P99	Surveillance of <i>Bacillus cereus</i> in Onigiri, Tobiko sushi and Sushi rice in Ladkrabang area	676
7	P114	Storage of ginger oil and its inhibition on <i>Colletotrichum musae</i>	682
8	P115	<i>Spirulina</i> as a protective agent on the survival of <i>Lactobacillus acidophilus</i> after freeze-drying	691
9	P130	Effects of microwave irradiation on bacteria in drinking water	696
10	P212	The feasibility study of <i>Centella asiatica</i> extract-loaded BSA nanoparticles production for antibacterial activity	703
11	P222	Antibacterial activity and chemical composition of essential oils from <i>Etlingera pavieana</i> (Pierre ex Gagnep.) R.M.Sm	707
12	P240	Fermentation of malva nut juicemixture by probiotic lactic acid bacteria	713
13	P73	Effects of various stresses on survivability of probiotic bacteria	719





## Substitution Effect of Mocaf (Modified Cassava Flour), Carrageenan and Seaweed on Properties of Analog Rice from Lesser Yam Tubers (*Dioscorea esculenta* L.)

Sri Winarti<sup>1)</sup>, Enny Karti Basuki Susiloningsih<sup>1)</sup> and Ramdan Hidayat<sup>2)</sup>

<sup>1</sup>Food Technology Program, Industrial Technology Faculty, University of Pembangunan Nasional “Veteran” Jawa Timur, Jl. Raya Rungkut Madya, Surabaya, Indonesia, 60294; Phone number: (031)8782179  
Corresponding author E-mail : [swin\\_tpupn@yahoo.com](mailto:swin_tpupn@yahoo.com)

<sup>2</sup>Faculty of Agriculture,, University of Pembangunan Nasional “Veteran” Jawa Timur, Jl. Raya Rungkut Madya, Surabaya, Indonesia, 60294

**ABSTRACT:** Research on diversification of food product from lesser yam tubers (*Dioscorea esculenta* L.) into analog rice was performed. This study was the innovation of new products to enhance the benefits of local food sources. The purpose of this study was to obtain the best formula of analog rice from lesser yam tubers, mocaf (modified cassava flour), carrageenan and seaweed. This research was conducted in two stages, the first stage was the making of lesser yam and mocaf flour. Making of mocaf carried out by fermentation using *Lactobacillus plantarum* FNCC 0027. The 2<sup>nd</sup> stage was the addition carrageenan or seaweed on the formulation of analog rice. The data from this research analysis using ANOVA (analysis of variant) and DMRT (duncan't multiple range teste). The results showed that the average yield of lesser yam flour 16.24%, mocaf flour 23.32% (fermentation 1 day); 23.23% (fermentation 2 days) and 22.73% (fermentation 3 days). The results of the studied known that the addition of 2% carrageenan in analog rice preferably consumers compared with addition of seaweed 2%. The analog rice with addition 2% carrageenan have characteristics: the levels of dietary fiber 14.28%; rehydration power 51.00% and expansion volume 136.67%, the value of consumer preferences to the taste of cooked rice was 3.67; texture 3.83; and smell 3.83. Sensory method used in research was hedonic scale scoring, with range value 1-5. 1 = not preference and 5= very preference by consumer.

**Keywords:** analog rice, lesser yam, carrageenan, seaweed, *Dioscorea esculenta* L.

### INTRODUCTION

Food and nutrition is one of the determinants of increased productivity and national competitiveness in the global arena. Supply, distribution and consumption of food with the amount, security and adequate nutritional quality must be guaranteed. Diet and desires of the community in various area in Indonesia so diverse that need

explore the potential of local food in each region so that people healthy and active life.

Indonesia is very rich variety of local food that is already exist and entrenched in society. Local food has a strategic role and it is tremendous potential, but currently there is a tendency for people ignoring local food sources and prefers imported food. The younger generation may be even don't know more local food in the area. The local food contains a lot of health benefits that are



not less than imported food and has a strategic role as well as many benefits for health.

Despite of the many benefits, unfortunately, people still seem less appreciate and realize the importance of developing local food seriously. As a result, people become accustomed with food imports. Base on regulation and food diversification of local resources, the Government of Indonesia hope that level of food consumption patterns must be diverse, nutritionally balanced, lawful and safe. Its needs strengthening and the participation of local governments in the development and implementation food diversification program based on local resources. One type of local tubers that need to be improved and empowered role is lesser yam tuber (*Dioscorea esculenta* L.).

Lesser yam is one kind of plant that grows in many parts of Indonesia, growing wild in the garden residents and in the forests. The tubers from this plant normally were used by community as source of carbohydrate alternative. Until now it is use still very limited.

The lesser yam plant advantages compared to other plants that can be grown on non-irrigated land, degraded land, without intensive farming and as intercrops (Gsianturi, 2003), the tubers contain inulin which has activity as a prebiotic (Winarti, *et al.*, 2011). Prebiotics are food components that can't be digested in the upper gastrointestinal tract, and can stimulate the selective growth and activity of beneficial bacteria in the digestive tract, such as bifidobacteria and lactobacilli, so as to improve the health of the host (Gibson, 2004; Pompei *et al.*, 2008; Gaggia *et al.*, 2010).

Development of analog rice from lesser yam tubers is one effort to new innovation to make beneficial of local food source that is relatively abundant in Indonesia, and providing alternative food that is cheap and nutritious, so it can be

beneficial to strengthening food self-sufficiency.

Analog rice is rice the result from innovations base on lesser yam flour, made in the form of granules look like with the rice. There are two methods for making the analog rice. Analog rice produced by granulation technology has a hard texture and appearance beyond rude so less preferred by consumers. Therefore, we developed analog rice by extrusion technology. This analog rice has better quality and more preferred by consumers.

Extrusion is a process where materials imposed by the force of the screw to flow in a narrow room that will undergo mixing and cooking as well. The main heat source the extrusion process from the conversion of mechanical (friction) is due to friction between the material and friction between materials with a screw. Work the thread also results in an accumulation of pressure in the extruder barrel, the material is forced out through the mold (die) that is small in size and return to normal pressure (atmospheric) instantaneously ie when the product through the die (Budi, *et al.*, 2013).

Previous research that has been done was development prebiotic and synbiotic food product based on lesser yam tuber (Winarti and Saputro, 2013) and the extraction of inulin from the *Dioscorea* spp. tubers as prebiotic component (Winarti *et al.*, 2011). But making analog rice has never been done from lesser yam tubers. Therefore it is necessary to do research on the innovation to making analog rice, the sensory test to know consumer preferences about analog rice from lesser yam tubers.

The research objective was to determine the effect of substitution mocaf flour (modified cassava flour), carrageenan and seaweed to quality and consumer preference analog rice from lesser yam tubers.

## MATERIAL AND METHODS

Materials: lesser yam tubers (*Dioscorea esculenta* L.) were obtained from the



Nganjuk region, East Java. Supporting materials include mocaf, corn starch, carrageenan and seaweed.

Material for analysis including alpha-amylase enzyme, phosphate buffer, pepsin enzyme, beta-amylase enzyme, ethanol, acetone and destilation water.

The equipment used in this research includes cabinet dryers, single screw extruder, disk mill, plastic strapping, vacuum sealer, soaking tub, strainers, cans and plastic tools.

**Method:** Stage 1, a) Preparation of lesser yam flour, b) Preparation of mocaf flour by fermentation using *Lactobacillus plantarum* FNCC 0027. Stage 2: Formulation analog rice, the best formulation from previous research (mocaf flour 15% and lesser yam flour 85%), mixed with seaweed or carrageenan at concentrations of 0%,1% and 2%. Coupled with other ingredients that are GMS (Glycerol Mono Stearate), skim milk, vegetable oil, and water then mixed until homogeneous. The flour mixture/dough was formed with an single extruder (length 1.8 m, with diameter die 2mm) at a temperature of 100°C to form granules of rice called analog rice. Analog rice then dried at 60°C, for 24 hours.

Hedonic scale scoring test was used to evaluation of sensory quality of cooked analog rice with 20 consumers. Range value in this test 1 until 5, 1= not preference and 5= very preference by consumer.

## RESULT AND DISCUSSION

Characteristics of lesser yam flour which used as raw material for making analog rice were presented in Table 1.

Starch content is one of the criteria for the quality of flour, both as food and non-food. Amylopectin content of the material is closely related to adhesion properties of these materials. The higher levels of amylopectin in rice cause the cook rice more closely/sticky. Starch tubers that have high amylopectin content also form the gel

stickier compared with low amylopectin when starch is heated.

Table 1 Characteristics of lesser yam flour

Component	Content (%)
Yield of powder	16.24 ± 0.238
Water	5.04 ± 0.059
Ash	0.99 ± 0.026
Starch	82.82 ± 0.14
Amylose	13.26 ± 0.08
Amylopectin	69.56 ± 0.12
Dietary fiber	10.77±0.03

Note: the average value of three replicates

MOCAF (Modified Cassava Flour) is a derivate product from cassava flour that uses the principle of modified cassava cells by lactic acid bacteria during fermentation (Subagio, 2007; Haryadi, 2011). The lactic acid bacteria used in this study was *Lactobacillus plantarum* FNCC 0047 obtained from the Center of Food and Nutrition Studies, Universitas Gadjah Mada, Yogyakarta. Characteristics of mocaf with different fermentation time were presented in Table 2.

Table 2 Characteristics of mocaf with different fermentation time.

Fermentation time (day)	Yield (%)	Water content (%)	Ash content (%)
0	23.03±0.12	4.82±0.01	1.02±0.04
1	23.32±0.14	4.90±0.00	0.99±0.01
2	23.23±0.09	4.84±0.01	0.82±0.00
3	22.73±0.08	4.73±0.02	0.76±0.00

Note: the average value of three replicates

Increasing fermentation time can reduce the yield, moisture and ash content in mocaf. That is due that the microbes were grow on cassava produce pectinolytic and cellulolytic enzymes that can destroy the cell walls of cassava such that happen destruction/perforation of the starch granules. The microbes also produce enzymes that hydrolyze starch into sugars and then convert it into organic acids,



especially lactic acid. The longer of fermentation time, starch granules are broken more so that simplify the process of evaporation of water during the drying mocaf, so the water levels began to decline. Decline of water levels in mocaf cause reduced the yield by weight.

During fermentation there are components of cassava soluble in water and occurs destruction of cellulose in cassava becomes soft texture and perforation on starch granules wall (Umar, 2013). The longer of fermentation time will be more cellulose wall ruptured, it is due the water content easier for evaporated, and declined the yield of mocaf.

The results of the analysis of starch granules mocaf using electron scanning microscope can be seen in Figure 1. The longer of fermentation time, the starch granules were damaged/broken more (indicated by arrows in Figure 1). This is due the microbes that grow on cassava will produce pectinolytic and cellulolytic enzymes that can destroy the cell walls of cassava granules. The microbes also produce enzymes that hydrolyze starch into sugars and then convert it into organic acids, especially lactic acid. This process will lead to changes in characteristics of the cassava flour include increasing the viscosity, gelation capability, rehydration power, and enhanced the solubility. Furthermore, the starch granules will hydrolysis to produce monosaccharide as raw material to produce organic acids. Compounds of this acid mixes with cassava flour cause produce sense and specific flavor that can mask the natural sense and flavor of cassava which not preferred by consumers. The flavor of mocaf become neutral to cover up the flavor of cassava to 70% (Subagio, 2007).

During fermentation can remove the color components such as pigments (especially the yellow cassava) and proteins that can cause a brown color when heating. The impact of removing the color components was the mocaf whiter than the

usual color of cassava flour and also smelled neutral (no musty smell typical). This process will produce flour that almost resembles the characteristics and quality of wheat flour, so that the product is suitable for replacing wheat flour for food industry.




Figure 1 SEM (scanning electron microscope) of mocaf (A) Fermentation time 0 day; (B) 1 day; (C) 2 days; (D) 3 days.

### Dietary Fiber of Analog Rice

The results showed that the higher addition of carrageenan or seaweed increase content of dietary fiber in the analog rice. This is because the main components of seaweed and carrageenan are soluble dietary fiber. Total dietary fiber is the total soluble fiber plus the total insoluble fiber (Marsono, 2002).




Figure 2 Total dietary fiber in analog rice  
Note: different letters following the values in each histogram indicates significantly different

### Rehydration Power





Rehydration power/water absorption of analog rice determined to know the ability of the rice to absorb water back when rice cooked. Water absorption can also be used to predict the shelf life of analog rice and appropriate packaging methods. Water absorption is influenced by several things, among others, long-chain, the number of polar groups or hydroxyl groups, the surface area of powder and water content (Hariyadi, 2011).

The results study showed that the higher addition of carrageenan/seaweed increased the rehydration power (Figure 3). Because carrageenan is a hydrocolloid compound that can form three-dimensional network when heated, so as to absorb and trap water well.




Figure 3 Rehydration power of analog rice. Note: different letters following the values in each histogram indicates significantly different

### Expansion Volume

The expansion volume of analog rice is the ability of rice to swell after steaming. Expansion volume has an important role to the quality of analog rice. The mechanism of expansion analog rice happened because rice absorbs the water through process of gelatinization of starch, starch properties can trap water to form a three-dimensional network.

The results from this research show that addition of carrageenan and seaweed can increase the expansion volume of the

analog rice (Figure 4). This is because the carrageenan and seaweed is a hydrocolloid compound that can trap high water when heated.

### Sensory Quality

The quality of food can be assessed in three ways: chemical, physical and sensory. Accepted or not the food products by consumers is determined by the quality factor especially sensory quality. Sensory properties are the nature of the starting materials by using human senses, namely the senses of sight, smell and taste.




Figure 4 Expansion volume of analog rice. Note: different letters following the values in each histogram indicates significantly different

The results showed that the highest score of consumer preference of taste, texture and color analog rice from lesser yam was the analog rice with addition carrageenan 2% (Table 3).

Table 3 The average of consumer preferred

Treatment	Teste	Texture	Colour
F0 (control)	3.00 <sup>ab</sup>	3.00 <sup>b</sup>	2.00 <sup>ab</sup>
Carragenan 1%	2.27 <sup>c</sup>	1.733 <sup>c</sup>	2.47 <sup>b</sup>
Carragenan 2%	3.40 <sup>a</sup>	3.89 <sup>a</sup>	3.80 <sup>a</sup>
Seaweed 1%	2.87 <sup>b</sup>	2.33 <sup>bc</sup>	2.53 <sup>b</sup>
Seaweed 2%	3.00 <sup>ab</sup>	2.53 <sup>bc</sup>	2.27 <sup>b</sup>

Note: different letters following the values indicates significantly different

Addition carrageenan 2% can increase chewy texture and fluffier than the addition of seaweed 2%. The addition of 2% carrageenan also provide color whiter than



seaweed 2%, so it is more preferred by consumers. Addition seaweed is tends to give texture sticky and wet. Analog rice from lesser yam tubers addition with seaweed or carrageenan presented in Figure 5A and 5B.




Figure 5 Analog Rice from Lesser Yam Tubers, (A) Addition with seaweed, (B) Addition with carrageenan

## CONCLUSIONS

The results of research indicated that the average of yield lesser yam flour 16.24%, mocaf flour 23.32% (fermentation 1 day); 23.23% (fermentation 2 days) and 22.73% (fermentation 3 days). The analog rice with addition 2% carrageenan have characteristics: the levels of dietary fiber 14.28%; rehydration power 51.00% and expansion volume 136.67%, the value of consumer preferences to the taste of cooked rice was 3.67; texture 3.83; and smell 3.83. Addition of carrageenan 2% in analog rice, preferably consumers compared with the addition of seaweed 2%.

## ACKNOWLEDGEMENTS

Acknowledgements to Ministry of Research and Technology Directorate General of Higher Education that has funded this research through Prime Research College of 2014 to 2016.

## REFERENCES

Budi, F.S., Hariyadi, P., Budijanto, S. and Syah, D. 2013. Extrusion Process Technology of Analog Rice. PANGAN, Vol. 22 No. 3: 263-274.

Gaggia, F., Mattarelli, P. and Biavati, B. 2010. Probiotics and prebiotics in animal feeding for safe food production. *International Journal of Food Microbiology* 141:515-528.

Gibson, G.R., 2004. Prebiotics. *Best Practice & Research Clinical Gastroenterology*. Vol.18, No.2, pp.287-298. (Chapter in book).

Gsianturi, 2003. Strengthening Food Security With Tubers. Article presented at National Food Industry, Technology Expert Association Food Indonesia.

Haryadi, 2011. Modified Cassava Flour Technology. *AGRITECH*, Vol. 31:2;86-92.

Marsono, Y. 2002. The glycemic index some tubers. Article presented at National Food Industry, Technology Expert Association Food Indonesia, Surabaya 10–11 October 2002.

Pompei, A., Cordisco, L., Raimondi, S., Amaretti, A. and Pagnoni, U.M. 2008. In vitro comparison of the prebiotic effect of two inulin-type fructans. *Anaerobe* 14:280-286.

Roberfroid, M.B., 2005. Introducing inulin-type fructans. *British Journal of Nutrition*, 93, Suppl.1, S13-S25

Setiawati, N.P., Santoso, J., Purwaningsih, S. 2014. THE CHARACTERISTICS OF ARTIFICIAL RICE WITH SEAWEED *Eucheuma cottonii* ADDITION AS A DIETARY FIBER SOURCE. *Journal of Tropical Marine Science and Technology*, Vol. 6, No.1:197-208.

Subagio, A. 2007. Industrialisasi Modified Cassava Flour (MOCAF) as raw material to Food Industry to support the diversification of the national food staple. Food Technology Faculty, Jember University.

Umar, M. 2013. MOCAF Flour as a substitute for wheat flour. <http://allinonenewfree.blogspot.com/2013>.

Winarti, S., E. Harmayani and R. Nurismanto. 2011. Ekstraksi of Inulin From Various Yam Tuber (*Dioscorea* spp.), article presented in AFC (Asian Food Conference), Bangkok 15-19 June 2011.

Winarti, S., dan E.A. Saputro. 2013. Characteristics of prebiotic flour from yam





tubers (*Dioscorea* spp.). Journal of Chemical Engineering, Vol.8, No.1: 17-21.

Winarti, S., Harmayani, E., Marsono, Y., and Pranoto, Y. 2013. Effect of inulin isolated from lesser yam (*Dioscorea esculenta* L.) on the growth of probiotics bacteria and SCFA formation during fermentation. International Research Journal of Microbiology (IRJM), Vol 4(2): 53-63.

